



An operational urban scale air quality forecast system

Mari Kauhaniemi
Ari Karppinen
Jari Härkönen
Juha Nikmo
Julius Vira
Marje Prank
Lasse Johansson
Jaakko Kukkonen

Background

- Aim
 - Information and planning tool for authorities
 - Information and warnings for public about high air pollution
- Requirements
 - Series of models from emission modelling and meteorological forecasting to dispersion of pollutants
 - Data retrieval and processing tools
 - Operational forecasting
 - Web interface for results

Urban air quality forecast system at FMI

•Includes



- Meteorological and air quality data retrieval
- Models for emission, chemistry, and dispersion (FORE, CAR-FMI)

•Limitations

- Considers only road traffic emissions


•Output

- 44 hour forecast four times a day (02, 08, 14, and 20 local time)
- NO₂, NO, CO, O₃, PM_{2.5}, and PM₁₀
- Domain is the Helsinki metropolitan area (40 km x 30 km; grid size 50-500 m)




Urban air quality forecast system at FMI

- Real-time results shown on website uaqfs.fmi.fi



Description of the system.





Example of model evaluation.

Dispersion and chemistry model (CAR-FMI)

- Contaminants in the Air from a Road (Härkönen, 2002)
- Traffic-originated pollution from an open road network
- Includes
 - Gaussian plume dispersion
 - Dry deposition of particles
 - Discrete parcel method for NO-O₃-NO₂ chemistry
 - Traffic-induced turbulence
- Evaluated e.g. by
 - NO₂: Levitin et al., 2005, Kukkonen et al., 2001, Kousa et al., 2001, Karppinen et al., 2000
 - PM_{2.5}: Kauhaniemi et al., 2008, Tiitta et al., 2002
 - PM_{2.5} and NO₂: Sokhi et al., 2008

Traffic emission modelling

- Exhaust emission factors for NO_x, CO, PM_{2.5}
 - National estimates
- Non-exhaust emission factors for PM₁₀
 - Road dust emission model (FORE)
- Traffic data (EMME-2 model)
 - Traffic volume
 - Travel speed (for exhausts)



Road dust emission model (FORE)

- Forecasting Of Road dust Emissions (Kauhaniemi et al., 2011)
- Based on PM emission model of SMHI (Omstedt et al., 2005)
- Considers
 - Moisture content of the road surface.
 - Particles from the wear of pavement due to tyres and traction sand.
- Not included
 - Emissions from brake, tyre, and clutch.
 - Dependencies of emissions on vehicle speed or fleet composition.
 - Influence of salting, dust binding, ploughing, and cleaning.
- Output
 - The emission factor of road wear and traction sand for all traffic in $\mu\text{g/veh/m}$.


Meteorological and regional background AQ data

- Meteorological forecasts from HIRLAM
 - High resolution limited area model
 - 54 h forecast
 - four times a day (at 00, 06, 12, 18 UTC)
- Background air quality forecasts from SILAM
 - System for integrated modelling of atmospheric composition
 - 72 h forecast
 - one time a day (at 02 UTC)




Correction of predicted background air quality

- The level of forecasted background concentrations is corrected with regional background air quality observations.





AQF start time t_0 = 02, 08, 14, and 20 local time


AQ measurement sites used for model evaluation




Results: Regional background by SILAM







Results: Hourly values summary







Results: Daily values summary



Results: Daily averaged PM2.5 and PM10



Results: Daily averaged NO₂ and O₃



Conclusions

- Accuracy of AQ forecasting system regarding daily values for t+24 forecast data:
 - PM2.5: fairly good
 - PM10, NO2 and O3: moderate
- PM2.5 usually somewhat over-predicted and NO2 and PM10 under-predicted.
- Correlation regarding daily values is better than for hourly values.

Challenges and further work

- Forecasting urban meteorology
 - Stability, precipitation, ...
- Forecasting of regional background concentrations
- Emission and traffic data up-to-date as possible
- PM10 modelling
 - Road dust emission model
- Other
 - Street canyons,... other emission sources e.g. small scale combustion?

References

- HSY, 2011. Photos of the study areas. The Helsinki Region Environmental Services Authority.
- Härkönen, J., 2002. Regulatory dispersion modelling of traffic-originated pollution. Finnish Meteorological Institute Contributions 38, Helsinki, University press, 103p.
- Karppinen A., Kukkonen J., Elolähde T., Konttinen M., Koskentalo T., 2000. A modelling system for predicting urban air pollution: comparison of model predictions with the data of an urban measurement network in Helsinki. Atm Env 34, 3735-3743.
- Kauhaniemi M., Karppinen A., Härkönen J., Kousa A., Alaviippola B., Koskentalo T., Aarnio P., Elolähde T. and Kukkonen J., 2008. Evaluation of a modelling system for predicting the concentrations of PM_{2.5} in an urban area. Atm Env 42, 4517-4529.
- Kauhaniemi M., Kukkonen J., Härkönen J., Nikmo J., Kangas L., Omstedt G., Ketzel M., Kousa A., Haakana M., Karppinen A. 2011. Evaluation of a road dust suspension model for predicting the concentrations of PM₁₀ in a street canyon. Atm Env 45, 3646-3654.
- Kousa A., Kukkonen J., Karppinen A., Aarnio P., Koskentalo T., 2002. Statistical and diagnostic evaluation of a new-generation urban dispersion modelling system against an extensive dataset in the Helsinki area. Atm Env 35, 4617-4628.
- Kukkonen J., Härkönen J., Walden J., Karppinen A., Lusa K., 2001. Evaluation of the CAR-FMI model against measurements near a major road. Atm Env 35, 949-960.
- Levitin J., Härkönen J., Kukkonen J., Nikmo J., 2005. Evaluation of the CALINE4 and CAR-FMI models against measurements near a major road. Atm Env 39, 4439-4452.
- Omstedt G., Bringfelt B., Johansson C., 2005. A model for vehicle-induced non-tailpipe emissions of particles along Swedish roads. Atm Env, 39, 6088-6097.
- Sokhi R.S., Mao H., Srimath S.T.G., Fan S., Kitwiroon N., Luhana L., Kukkonen J., Haakana M., Karppinen A., van den Hout K.D., Boulter P., McCrae I.S., Larssen S., Gjerstad K.I., San José R., Bartzis J., Neofytou P., van den Breemer P., Neville S., Kousa A., Cortes B.M., Myrtveit I., 2008. An integrated multi-model approach for air quality assessment: Development and evaluation of the OSCAR Air Quality Assessment System. Env Mod & Soft 23, 268-281.
- Tiitta P., Raunemaa T., Tissari J., Yli-Tuomi T. Leskinen A., Kukkonen J., Härkönen J., Karppinen A., 2002. Measurements and modelling of PM_{2.5} concentrations near a major road in Kuopio, Finland. Atm Env 36, 4057-4068.




Thank you!


System performance study

- 24 h forecast of PM_{2.5}, PM₁₀, NO₂, O₃ concentrations
- Values $t+24 \dots t+30$ picked from each result file and combined together for one time series.
- Time period: 1 June 2013 to 31 May 2014

44 h result data



*Forecast run start time
 $t_0 = 02, 08, 14, \text{ and } 20$
local time*







*Combined data,
i.e., study data*






Results: Hourly values



Comparison with previous studies with observed bg and met data

NOTE! Not directly comparable with this study.



- **VALLILA NO2 (1996, 1997):** Evaluation of modelling system with observed met. (MPP-FMI) and regional background data, emi. from stationary sources included (Kousa et al., 2001)
- **KALLIO PM2.5 (2002):** Modelling system with observed met. (MPP-FMI) and regional background data, non exh. emi. included by coefficient (Kauhaniemi et al., 2008)
- **VALLILA PM10 (2004):** Modelling system with observed met. (MPP-FMI) and regional background data, road dust emi. Included